

Cell Cycle Arrest and Cell Death Are Controlled by p53-dependent and p53-independent Mechanisms in Tsg101-deficient Cells*

Received for publication, January 14, 2004, and in revised form, May 5, 2004
Published, JBC Papers in Press, June 21, 2004, DOI 10.1074/jbc.M400408200

Marissa J. Carstens^{‡§}, Andrea Krempler^{‡¶}, Aleata A. Triplett[‡], Maarten van Lohuizen^{**},
and Kay-Uwe Wagner^{‡ ‡‡}

From the [‡]Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, Nebraska Medical Center, Omaha, Nebraska 68198-6805 and ^{**}The Netherlands Cancer Institute, Department of Molecular Genetics, H5, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

Our previous studies have shown that cells conditionally deficient in Tsg101 arrested at the G₁/S cell cycle checkpoint and died. We created a series of Tsg101 conditional knock-out cell lines that lack p53, p21^{Cip1}, or p19^{Arf} to determine the involvement of the Mdm2-p53 circuit as a regulator for G₁/S progression and cell death. In this new report we show that the cell cycle arrest in Tsg101-deficient cells is p53-dependent, but a null mutation of the p53 gene is unable to maintain cell survival. The deletion of the *Cdkn1a* gene in Tsg101 conditional knock-out cells resulted in G₁/S progression, suggesting that the p53-dependent G₁ arrest in the *Tsg101* knock-out is mediated by p21^{Cip1}. The Cre-mediated excision of *Tsg101* in immortalized fibroblasts that lack p19^{Arf} seemed not to alter the ability of Mdm2 to sequester p53, and the p21-mediated G₁ arrest was not restored. Based on these findings, we propose that the p21-dependent cell cycle arrest in Tsg101-deficient cells is an indirect consequence of cellular stress and not caused by a direct effect of Tsg101 on Mdm2 function as previously suggested. Finally, the deletion of *Tsg101* from primary tumor cells that express mutant p53 and that lack p21^{Cip1} expression results in cell death, suggesting that additional transforming mutations during tumorigenesis do not affect the important role of Tsg101 for cell survival.

The tumor susceptibility gene 101 (*Tsg101*) encodes a multidomain protein that mediates a variety of biological functions. Some of these functions have been postulated from the predicted protein structure, its intracellular localization, the identification of Tsg101-binding proteins, and very recently, the generation of Tsg101-deficient mouse models. These functions include a role in ubiquitination (1, 2), transcriptional regulation (3, 4), endosomal trafficking (5–8), proliferation (9–12), and cell survival (12, 13).

* This work was supported by NCI, National Institutes of Health Public Health Service Grant CA93797 (to K.-U. W.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ Recipient of a stipend from Department of Defense Grant DAMD17-00-1-0361.

¶ Supported by Deutsche Forschungsgemeinschaft Grant DFG, KR 2107/1-1.

|| Present address: University of the Saarland, Medical Biochemistry and Molecular Biology, Bldg. 44, Homburg, 66424, Germany.

‡‡ To whom correspondence should be addressed: Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, 986805 Nebraska Medical Center, Rm. 8009, Omaha, NE 68198-6805. Tel.: 402-559-3288; Fax: 402-559-4651; E-mail: kuwagner@unmc.edu.

The C-terminal coiled-coil domain of Tsg101, formally known as CC2, was isolated first in a yeast two-hybrid screen with the cell growth regulating protein Stathmin (14). This region is a potential co-repressor that is able to modulate the transcriptional activity of steroid receptors (3, 4, 15). Furthermore, Tsg101 possesses a proline-rich sequence known to exist in activation domains of transcription factors (16). The N-terminal region of Tsg101 (UEV domain) is similar to the catalytic domain of ubiquitin-conjugating (E2) enzymes. Because Tsg101 lacks a key cysteine residue found in authentic E2 enzymes, it has been postulated that this protein might serve as a negative regulator for the ubiquitin-mediated degradation of other proteins (1, 2). Suggested targets of important factors for cell cycle regulation whose stability and function were affected by Tsg101 are Mdm2 and p21^{Cip1} (17, 18).

New insights about the biological role of Tsg101 *in vivo* were obtained from genetically engineered mice that lack Tsg101 completely (11, 13) or in selected cell types (12, 13). The deletion of the promoter and the first coding exon of *Tsg101* resulted in embryonic lethality around implantation (13). Tsg101-deficient embryos lacking exons 8 and 9 die around day 6.5 of gestation due to a defect in cell proliferation and mesoderm formation (11). The WAP¹-Cre-mediated deletion of Tsg101 from differentiating mammary epithelial cells (conditional knock-out) in adult females resulted in increased cell death, impaired mammary development, and a lactation-deficient phenotype (13). The excision of the *Tsg101* gene in primary cultures of mouse embryonic fibroblasts (MEFs) or mammary epithelial cells (MECs) revealed that Tsg101-deficient cells arrested at the G₁/S transitional phase of the cell cycle before they underwent cell death (12, 13). In contrast to previous reports (16, 19), neither haplo-insufficiency of *Tsg101* nor the deletion of both *Tsg101* alleles in the conditional knock-out models (*in vitro* and *in vivo*) resulted in neoplastic transformation, suggesting that a null mutation of *Tsg101* is not an initiating event for tumorigenesis (11–13).

Cell cycle arrest at the G₁/S checkpoint and cell death were two major phenomena that we observed in the Tsg101 conditional knock-out model (12, 13). The analysis of crucial regulators of the cell cycle revealed that Tsg101 deficiency resulted in growth arrest through the inactivation of cyclin-dependent kinase 2 (Cdk2). Consequently, DNA replication was not initi-

¹ The abbreviations used are: WAP, whey acidic protein; Cre, site-specific recombinase in bacteriophage P1 (catalyzes recombination between loxP sites (loxP, locus of X-ing over)); Mdm2, transformed mouse 3T3 double minute 2 protein; MEF, mouse embryonic fibroblast; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Cdk2, cyclin-dependent kinase 2; MEC, mammary epithelial cell.



A**B**

FIG. 1. p53 deficiency does not rescue cell death in the Tsg101 conditional knock-out. *A*, *in vitro* analysis. An MTT color assay was performed to determine the growth rates of pBabe and pBabe-Cre-infected *Tsg101*^{fl/fl} MEFs, which lack one or two alleles of the *p53* gene (*p53*^{+/-} and *p53*^{-/-}) and their wild type controls (*p53*^{+/+}). The $A_{570\text{ nm}}$ values that correspond to total cell numbers decrease in the Tsg101 knock-out (*Tsg101*^{fl/fl} pBabe-Cre) between 4 and 7 days after retroviral Cre infection regardless of the *p53* mutation status. In contrast, the number of cells in the pBabe-infected controls increase steadily. *Error bars* correspond to S.D. *B*, *in vivo* analysis. Whole mounts of carmine alum-stained mammary glands from tissue-specific Tsg101 knock-out mice (WAP-Cre *Tsg101*^{fl/fl} *p53*^{+/+}), females that are conditionally deficient in both, Tsg101 and p53 (WAP-Cre *Tsg101*^{fl/fl} *p53*^{-/-}), and their controls (*Tsg101*^{fl/fl} *p53*^{+/+}). Mammary glands were taken, fixed, and stained for several hours post-partum (magnification 40 \times). Note that alveolargenesis is severely impaired in the Tsg101 conditional knock-outs regardless of whether p53 is expressed or not.

ated in Tsg101-deficient cells before they died (12). A crucial regulatory mechanism, which is commonly associated with G₁ arrest and cell death, is the p19^{Arf}-Mdm2-p53 tumor-surveillance or stress-response pathway (20). The goal of this study was to further discriminate the p53-dependent and p53-independent mechanisms that lead to either cell cycle arrest or cell death as a consequence of Tsg101 deficiency. We now show that the G₁ arrest in Tsg101-deficient cells is dependent upon the presence of functional p53 and its downstream mediator p21^{Cip1}. Our findings suggest that, in contrast to keratinocytes (18), Tsg101 is not required for p21^{Cip1} protein stability and function in proliferating fibroblasts. The absence of either p53 (and, therefore, p21) or p21^{Cip1} alone is, however, unable to sustain cell survival. In addition to these findings *in vitro*, a null mutation of the *p53* gene does not restore normal mammary glandogenesis and lactation in Tsg101 mammary-specific knock-

out mice (WAP-Cre *Tsg101*^{fl/fl} *p53*^{-/-}). To address the proposed role of Tsg101 as a crucial regulator for Mdm2 function (17), we have generated a conditional double-knock-out of *Tsg101* and *Cdkn2a*. The deletion of Tsg101 in immortalized fibroblasts that lack expression of p19^{Arf} does not alter Mdm2 function and does not restore the p21-mediated G₁ arrest. In summary, our data do not support a biologically relevant function of Tsg101 as a stabilizer for Mdm2. Therefore, we propose that the p21-mediated induction of the G₁ arrest might be an indirect consequence of Tsg101 deficiency due to cellular dysfunction and stress. Finally, we show that the deletion of Tsg101 from tumorigenic cells that express mutant p53 and that lack p21^{Cip1} expression results in cell death. This observation suggests that additional, transforming mutations during tumorigenesis do not affect the important role of Tsg101 for cell survival.






FIG. 2. A p53 null mutation restores G₁/S progression and the activity of Cdk2 in Tsg101-deficient cells. *A*, flow cytometric analysis of the DNA content of viable Tsg101-deficient MEFs (*Tsg101*^{fl/fl}, pBabe-Cre) that lack one or two alleles of the *p53* gene (*p53*^{+/-} and *p53*^{-/-}) and their uninfected controls (*Tsg101*^{fl/fl} *p53*^{-/-}). The sub-G₁ population of apoptotic cells was gated out in this assay. Note that the relative number of cells in S phase is reduced only in *p53* heterozygous knock-out cells lacking Tsg101 but not in Tsg101-deficient MEFs carrying two mutant *p53* alleles. *B*, Western blot analysis of cyclin A (S-phase cyclin) and regulators of G₁/S progression (p19^{Arf} and p21^{Cip1}) in Tsg101-deficient MEFs that lack one or two copies of *p53* and their uninfected controls. Note that cyclin A2 was markedly down-regulated in Tsg101-deficient MEFs that carry at least one functional allele of *p53*. Immortal *p53*/p21^{Cip1}-deficient MEFs express high levels of cyclin A2 and p19^{Arf} regardless of the *Tsg101* mutation status. *C*, Cdk2 activity assay using histone H1 as a substrate for phosphorylation. Only *Tsg101* knock-out MEFs that carry at least one functional *p53* allele exhibit a reduced activity of Cdk2. The complete deletion of *p53* and, subsequently, the absence of the Cdk2 inhibitor p21^{Cip1} restore a normal activity of Cdk2 in Tsg101-deficient cells.

A**B**

Genotype	G0/G1	S-Phase	G2/M
<i>Tsg101^{fl/fl} p21^{-/-} pBabe</i>	39.7 (±6.7) %	20.8 (±2.4) %	39.5 (±5.2) %
<i>Tsg101^{fl/+} p21^{-/-} pBabe-Cre</i>	59.6 (±9.1) %	18.4 (±3.5) %	22.0 (±9.5) %
<i>Tsg101^{fl/fl} p21^{-/-} pBabe-Cre</i>	45.7 (±7.9) %	22.4 (±3.0) %	31.9 (±7.1) %

FIG. 3. p21^{Cip1} is a mediator of the cell cycle block at the G₁ checkpoint in Tsg101 conditional knock-out cells. *A*, Western blot analysis of Tsg101 and p21^{Cip1} to monitor effective Cre-mediated excision and down-regulation of Tsg101 in MEFs lacking one or two copies of the *Cdkn1a* gene and their controls. *B*, flow cytometric analysis of the DNA content in cells doubly deficient in Tsg101 and p21^{Cip1} (*Tsg101^{fl/fl} p21^{-/-} pBabe-Cre*) and their controls. The sub-G₁ population of apoptotic cells was gated out in this assay. Note that the relative number of cells in S phase is normal in Tsg101-deficient MEFs carrying two mutant *Cdkn1a* alleles compared with their controls ($p > 0.05$).

EXPERIMENTAL PROCEDURES

Mouse Models, Genotyping Protocols, and Whole Mount Analysis of Mammary Glands—The PCR protocol for genotyping the whey acidic protein (WAP)-Cre mice (TgN(Wap-cre)11738Mam) as well as the generation and phenotypic characterization of *Tsg101* conditional knock-out mice (*Tsg101^{tm1Kuw}*) have been described earlier (13, 21). Mutant mice with a targeted deletion of the *p53* gene (*Trp53^{tm1Brd}*) (22) were purchased from Taconic Farms, Inc. *Cdkn1a* (23) and *Cdkn2a* (24) knock-out mice (*Cdkn1a^{tm1Tyj}* and *Cdkn2a^{tm1Rdp}*, respectively) were obtained from Jackson Laboratories and the repository of the Mouse Model for Human Cancer Consortium (MMHCC). Athymic nude mice (NCR strain, NCI) were used for transplantation studies. The preparation and staining of mammary gland whole mounts were described previously (13, 25). All animals used in the studies were

treated humanely and in accordance with federal guidelines and institutional policies.

Primary Cell Cultures and Retroviral Expression Vectors—MEFs from 13.5- or 14.5-day-old *Tsg101^{fl/fl}*, *Tsg101^{fl/fl} p53^{-/-}*, *Tsg101^{fl/fl} Cdkn1a^{-/-}*, or *Tsg101^{fl/fl} Cdkn2a^{-/-}* embryos and their littermate controls were explanted and maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, 2 mM glutamine, 0.1 mM nonessential amino acids, 10 μg/ml gentamycin, 100 units/ml penicillin, and 100 μg/ml streptomycin (Invitrogen). Cells at passages 2–4 were plated at a density of 4×10^5 cells/10-cm culture dish and infected with the pBabe and pBabe-Cre constructs. The generation of these retroviral vectors was published previously (12, 13). Forty-eight hours after infection cells were selected in complete medium containing 7 μg/ml puromycin (Sigma) or 200 μg/ml hygromycin B (Invitrogen).

Cell Cycle Analysis and 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide (MTT) Assay—Approximately 10^6 MEFs were harvested 4 days after puromycin selection to determine cell cycle progression using a flow cytometric analysis. Cells were washed in $1\times$ phosphate-buffered saline and fixed in ice-cold 70% ethanol for 30 min. After an additional washing step, cells were stained overnight with propidium iodide as described previously (26). The stained MEFs were analyzed in a FACScalibur (BD Biosciences) flow cytometer. The software packages CELLquest (BD Biosciences) and Modfit LT (Verity) were used for data acquisition. The cell cycle analysis was repeated three or four times for each experimental setting, and a *t* test was performed to validate statistically significant differences between the various double knock-out cell lines and their controls. An MTT growth assay was performed as described earlier (27) to determine the growth properties of *Tsg101/p53* double knock-out cells. The MTT was obtained from Sigma. 2×10^4 cells of each genotype were seeded in triplicates in a 96-well microtiter plate. Absorbance was measured at 570 nm with an Elx 808 (Bio-Tek Instruments) enzyme-linked immunosorbent assay reader.

Western Blot Analysis and Cdk2 Kinase Assay—MEFs were pelleted and lysed on wet ice for 30 min in $1\times$ phosphate-buffered saline, 1% Nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS, 1 mM phenylmethylsulfonyl fluoride, 0.4 units/ml aprotinin, 1 mM NaF, 0.1 mM sodium orthovanadate. Protein was quantified using a Bradford assay (Pierce) according to the manufacturer's protocol. Approximately 50–100 μ g of protein per lane was resolved on a 4–20% SDS-PAGE gradient gel and blotted onto polyvinylidene fluoride membranes (Invitrogen). The membranes were blocked for 1 h in $1\times$ Tris-buffered saline, 0.1% Tween 20, and 5% dry milk. Subsequently, membranes were incubated with primary antibodies in blocking buffer at 4 °C overnight, washed 3 times for 15 min in washing buffer ($1\times$ Tris-buffered saline, 0.1% Tween 20), and incubated for 1 h at room temperature with horseradish peroxidase-conjugated secondary antibodies in blocking buffer. Membranes were washed again 3 times in washing buffer and once for 15 min in $1\times$ Tris-buffered saline without Tween 20. Protein bands were detected using the ECL chemiluminescence kit for Western blot analysis (Amersham Biosciences) according to the manufacturer's protocol. Membranes were stripped using 0.2 M NaOH for consecutive detection of various proteins. The following antibodies were used in this study: α -Tsg101 (C-2), α -cyclin B1 (M-20), α -cyclin A2 (C-19), α -p16^{Ink4a} (M-156), and α -ActB (I-19) from Santa Cruz Biotechnology as well as α -p21^{Cip1} (SX118) from Pharmingen and α -p19^{Arf} (Ab-1) from Oncogene at a 1:1000 dilution. Horseradish peroxidase-conjugated secondary antibodies were purchased from Santa Cruz Biotechnology and used at a 1:1000 dilution. A Cdk2 kinase assay was performed using the α -Cdk2 (M-2) antibody from Santa Cruz Biotechnology and Histone H1 (1 μ g/ μ l) from Sigma.

RESULTS

A Null Mutation of p53 Does Not Rescue the Deleterious Phenotype Caused by Tsg101 Deficiency in Vitro and in Vivo—The Cre-mediated excision of both *Tsg101* floxed alleles from proliferating primary fibroblasts (MEFs) and MECs resulted in cell cycle arrest and cell death (12, 13). The activation of the p19^{Arf}-Mdm2-p53 stress-response pathway is frequently associated with both phenomena. To further discriminate the p53-dependent and p53-independent mechanisms that lead to either cell cycle arrest or cell death as a consequence of Tsg101 deficiency, we derived MEFs from Tsg101 conditional knock-out mice (*Tsg101*^{fl/fl}) that carry a targeted null mutation of the *p53* gene (22). Cells treated with a control virus (pBabe) or a retrovirus expressing Cre recombinase (pBabe-Cre) were grown for 7 days after infection to quantitatively examine the growth of Tsg101 knock-out cells in the presence or absence of p53. The multiplication of Tsg101-deficient MEFs and their controls were determined using an MTT assay between days four through seven post-infection when optimal puromycin selection, expression of Cre, and excision of *Tsg101* had been achieved (Fig. 1A). The Cre-mediated deletion of *Tsg101* resulted in a severe growth inhibition and reduction in the number of viable cells. More importantly, we did not observe a significant modification of growth properties or prolonged cell survival in the presence or absence of one or both copies of the *p53* gene. Our data suggested that a *p53* null mutation was




FIG. 4. The p19^{Arf}-p53 tumor surveillance and stress response pathway and its proposed interaction with Tsg101. *A*, Mdm2 is negatively regulated by p19^{Arf}. As a ubiquitin ligase, Mdm2 regulates the ubiquitin-mediated degradation of the tumor suppressor protein p53. Therefore, p19^{Arf}-deficient cells are immortal and lack expression of p53 and p21^{Cip1}. In normal cells, p21^{Cip1} is transcriptionally regulated by p53. It has been suggested recently that Tsg101 is a positive regulator for Mdm2 protein levels and function. Therefore, the deletion of the *Tsg101* gene (*right panel*) should destabilize Mdm2 and induce a p53/p21^{Cip1}-mediated G₁ arrest in cells lacking p19^{Arf}. *B*, experimental design to study the proposed interaction of Tsg101 with Mdm2/p53 in immortalized cells lacking p19^{Arf}. The Cre-mediated deletion of *Tsg101* in *Cdkn2a*^{-/-} immortalized cells should have a biologically relevant effect on Mdm2 and p53 protein levels and restore the cell cycle arrest.

neither able to rescue growth inhibition nor significantly modify the rate of cell death of Tsg101-deficient primary fibroblasts. We have shown in a control experiment that the expression of Cre recombinase can be excluded as a cause or possible modifier of this phenotype. Tsg101 wild type cells (*Tsg101*^{+/+}) infected with the retroviral pBabe-Cre vector did not exhibit growth retardation in an MTT assay. In addition, we were able to rescue the deleterious phenotype of *Tsg101*^{-/-} cells through the expression of exogenous, hemagglutinin-tagged Tsg101 from a retroviral vector (12).

To verify these initial findings in a different cell type and *in vivo*, we bred two mutant *p53* alleles into mammary-specific Tsg101 conditional knock-out mice (WAP-Cre *Tsg101*^{fl/fl} *p53*^{-/-}). The expression of Cre under the WAP promoter is largely confined to differentiating mammary epithelial cells in late-pregnant and lactating females (21, 25). In contrast to littermate controls (*Tsg101*^{fl/fl} *p53*^{+/+}), the WAP-Cre-mediated deletion of *Tsg101* in a *p53* wild type background (WAP-Cre *Tsg101*^{fl/fl} *p53*^{+/+}) resulted in impaired mammary development at the onset of lactation (Fig. 1B). The introduction of two mutant copies of the *p53* gene into the Tsg101 conditional knock-out (WAP-Cre *Tsg101*^{fl/fl} *p53*^{-/-}) was unable to rescue impaired mammary development caused by the WAP-Cre-mediated excision of the two floxed *Tsg101* alleles. Therefore, our observations in mutant female mice were consistent with the cell culture studies on *Tsg101/p53* double mutant MEFs. Taken together, the outcome of both studies (*in vitro* and *in vivo*) suggests that the inability of a *p53* null mutation to rescue a Tsg101-deficient phenotype is neither a cell culture phenomenon nor is it caused




FIG. 5. Cell cycle regulation in immortal cells that lack *Cdkn2a* and *Tsg101*. A, the tumorigenic, Tsg101-antisense-expressing SL6 cell line and its parental mouse 3T3 fibroblast cell line lack expression of p19^{Arf} and p16^{Ink4a}. *Cdkn2a*-deficient cells were used as negative controls, and immortal *p53*^{-/-} MEFs served as positive controls for p19^{Arf} and p16^{Ink4a} protein expression. B, Western blot analysis of Tsg101 and Cdkn2a to monitor effective Cre-mediated down-regulation of Tsg101 in MEFs lacking one or two copies of the *Cdkn2a* gene and their controls. The steady-state levels of Mdm2 did not decrease and were slightly elevated in cells lacking Tsg101. C, flow cytometric analysis of the DNA content in cells doubly deficient in Tsg101 and p19^{Arf} (*Tsg101*^{fl/fl} *Cdkn2a*^{-/-} pBabe-Cre) and their controls. The sub-G₁ population of apoptotic cells was gated out in this assay. P19^{Arf}-deficient cells deprived of essential growth factors (*Tsg101*^{fl/fl} *Cdkn2a*^{-/-} 0.1% fetal bovine serum (FBS)) served as an additional positive control to monitor the accumulation of cells at G₁ and the relative decrease in the number of cells at the S-phase of the cell cycle. Note that the deletion of *Tsg101* in a p19^{Arf}-deficient background does not cause a cell cycle arrest, whereas growth factor withdrawal from these cells results in a sharp reduction of cells in S phase.

Genotype	G0/G1	S-Phase	G2/M
<i>Tsg101</i> ^{fl/fl} <i>Cdkn2a</i> ^{-/-} pBabe	56.2 (±8.4) %	26.9 (±2.7) %*	16.9 (±7.3) %
<i>Tsg101</i> ^{fl/fl} <i>Cdkn2a</i> ^{-/-} pBabe-Cre	56.7 (±7.6) %	20.6 (±3.2) %*	22.7 (±9.5) %
<i>Tsg101</i> ^{fl/fl} <i>Cdkn2a</i> ^{-/-} 0.1% FBS	85.6 (±1.6) %	6.1 (±1.2) %	8.3 (±2.6) %

* P>0.05

by a possible difference in p53 function between different cell types of mesodermal and ectodermal origin (*i.e.* fibroblasts and mammary epithelial cells).

Deletion of *p53* in *Tsg101*-deficient Cells Restores the Activity of *Cdk2* and Progression into the S Phase—Despite p53-independent mechanisms that trigger the dominant phenotype (*i.e.* lethality) of *Tsg101*^{-/-} cells, we postulated that deficiency of

p53 might modify or even eliminate the cell cycle arrest in Tsg101-deficient cells before they die. To address this hypothesis, we analyzed the cell cycle progression of Tsg101 null cells lacking one or two copies of the *p53* gene. The Cre-mediated deletion of *Tsg101* in MEFs with a heterozygous null mutation of *p53* (*Tsg101*^{fl/fl} *p53*^{+/-} pBabe-Cre) leads to a significant reduction in the number of cells in S phase (Fig. 2A) (*p* < 0.01;




FIG. 6. Tsg101 deficiency causes cell death of tumorigenic cells. *A*, experimental design to address whether *Tsg101* is dispensable or essential for the survival of neoplastic cells. *B*, the subcutaneous injection of immortal *Tsg101^{fl/fl}* 3T3 fibroblasts expressing mutant p53 into *whn^{-/-}* nude mice (NCr strain) form solid tumors. *C*, hematoxylin and eosin staining of histological sections from solid tumors (magnification 100 \times). Note that these lesions are highly vascularized, and tumor cells invade into adjacent normal tissues. V, vasculature; NT, necrotic tumor tissue; MG, mammary gland; M, muscle. *D*, explanted tumor cells carrying two floxed alleles of *Tsg101* (*Tsg101^{fl/fl}*) infected with pBabe (control) or the pBabe-Cre retroviral vector to excise both copies of *Tsg101* (magnification 200 \times). Note that additional, transforming mutations during neoplastic transformation are incapable of rescuing lethality caused by *Tsg101* deficiency. *E*, MTT color assay to quantify the lack of cell growth of tumor cells lacking *Tsg101* compared with their controls. Error bars correspond to S.D.

t test), a decrease of the cyclin A2 protein level (Fig. 2B, *third lane*), and the inactivation of the Cdk2 (Fig. 2C). Consequently, the inhibition of only one *p53* allele had no effect on impaired G₁/S progression of *Tsg101*^{-/-} cells, and the consequences of *Tsg101* deficiency were identical to cells with two functional copies of *p53* (12). In contrast, the deletion of both *p53* alleles in *Tsg101* conditional knock-out MEFs (*Tsg101*^{fl/fl} *p53*^{-/-} pBabe-Cre) re-established a normal G₁/S progression (Fig. 2A) ($p < 0.02$; *t* test), increased the expression of cyclins A2 (Fig. 2B), and restored the activation of Cdk2 (Fig. 2C). In summary, these observations confirm our previously stated hypothesis that *p53* deficiency is able to lift the G₁ cell cycle arrest and to revert part of the complex phenotype of *Tsg101* knock-out cells despite its inability to sustain cell survival.

P21^{Cip1} Is a Mediator of the G₁ Cell Cycle Arrest in *Tsg101* Conditional Knock-out Cells—The activity of Cdk2 in complex with cyclin E is negatively regulated by the cell cycle inhibitor p21^{Cip1} whose expression level is dependent upon the transcriptional activation of *p53* (28–30). Therefore, we hypothesized that p21^{Cip1} might be a component of the *p53*-dependent cell cycle arrest in response to *Tsg101* deficiency. To address this issue, we bred *Cdkn1a*^{-/-} mice (23) into the *Tsg101*-floxed background (*Tsg101*^{fl/fl} *p21*^{-/-}) to generate MEFs that lack expression of p21 and *Tsg101* after infection with pBabe-Cre and selection with puromycin (Fig. 3A). The infection of p21^{Cip1}-deficient cells with the control vector pBabe and selection with puromycin had little or no consequences on the viability and multiplication of primary cells. The Cre-mediated excision of only one floxed allele of *Tsg101* (*Tsg101*^{fl/+} *p21*^{-/-} pBabe-Cre) also did not affect the survival and G₁/S progression of p21^{Cip1}-deficient MEFs as determined by flow cytometry (Fig. 3B). The deletion of two floxed alleles of *Tsg101* (*Tsg101*^{fl/fl} *p21*^{-/-} pBabe-Cre) progressively led to the death of *Tsg101*-deficient MEFs within 7 days after infection with the retroviral Cre vector (data not shown). Like the deletion or functional inhibition of *p53*, a null mutation of the *Cdkn1a* gene was, however, able to restore the G₁/S progression in cells lacking *Tsg101* (Fig. 3B). The average number of cells in S phase was not significantly different between the double knock-out cells and their controls ($p > 0.05$). The relative increase in a subset of cells at the G₂/M phase is probably caused by the puromycin selection since we observed the same phenomenon when these cells were infected with the pBabe vector control. Collectively our findings confirmed the working hypothesis, which predicted that p21^{Cip1} is a mediator of the G₁ cell cycle arrest in *Tsg101*-deficient cells.

The Deletion of Tsg101 in Immortalized Fibroblasts That Lack p19^{Arf} Does Not Alter the Ability of Mdm2 to Sequester *p53*—Several reports propose a potential role for *Tsg101* as an important negative regulator for the ubiquitin-mediated turnover of Mdm2 and p21^{Cip1} in various cell lines and in differentiating primary keratinocytes (11, 17, 18). In addition, Oh *et al.* (28) suggest that *Tsg101* is critical for the p21-mediated inhibition of cyclin-Cdk complexes in proliferating keratinocytes (18). The cell cycle analysis in double knock-out MEFs lacking *Tsg101* and *Cdkn1a* revealed that p21^{Cip1} is important for the G₁ arrest (see the previous paragraph). This observation suggests that a knock-out of *Tsg101* does not impair the stability, ubiquitin-mediated proteolysis, or function of p21^{Cip1} in proliferating fibroblasts. To address the proposed function of *Tsg101* as a stabilizer for Mdm2, we designed a new set of experiments illustrated in Fig. 4. Thus far, we determined the role of *p53* and p21^{Cip1} as regulators for the G₁/S progression in *Tsg101*-deficient cells. Both proteins act downstream of Mdm2 (20). In this new experimental design we planned to immortalize primary MEFs upstream of Mdm2 through deletion of the

Cdkn2a gene (Fig. 4A, *left*). The *Cdkn2a* locus encodes p19^{Arf}, which is known to negatively regulate Mdm2 (20). Therefore, p19^{Arf}-deficient immortalized cells also lack expression of *p53* and p21^{Cip1}. The level of unrestrained Mdm2 protein for the sequestration of *p53* is crucial in immortal p19^{Arf}-null cells.² This causal relationship is probably best illustrated by the fact that the deletion of *p19*^{Arf} alone has no influence on the survival of *Mdm2* mutant mice, whereas *p53/Mdm2* double knock-outs (31, 32) or *p19*^{Arf}/*Mdm2/p53* triple mutant mice are viable (33). According to this mechanism, the destabilization and down-regulation of Mdm2 as a consequence of *Tsg101* deficiency (Fig. 4A, *right*) should result in an accumulation of *p53* and p21^{Cip1}. In turn, the up-regulation of *p53* and p21^{Cip1} should inhibit cell cycle progression in *Cdkn2a* mutant cells. In conclusion, if *Tsg101* is essential for the stability and function of Mdm2 as previously suggested (17), then its deletion in immortalized MEFs lacking p19^{Arf} should impair G₁/S progression in a very similar fashion as in *Tsg101*-deficient nonimmortalized cells with normal *p53* and p21^{Cip1} levels.

To generate a conditional double-knock-out of *Tsg101* and *Cdkn2a*, we decided to cross *Tsg101*-floxed animals with *Cdkn2a* knock-out mice (24) that lack both tumor susceptibility proteins (p19^{Arf} and p16^{Ink4a}) encoded by this locus. Our decision was based on the fact that *Tsg101* and its proposed biological functions were first described in murine 3T3 fibroblasts and their derived tumorigenic SL6 cells (16). These cell lines lack p16^{Ink4a} in addition to p19^{Arf} as determined in a preliminary study (Fig. 5A). To generate primary *Tsg101/p19*^{Arf} double knock-out cells, we infected MEFs carrying two floxed alleles of *Tsg101* in addition to two null alleles of *Cdkn2a* (*Tsg101*^{fl/fl} *Cdkn2a*^{-/-}) with the pBabe-Cre virus (Fig. 5B). *Tsg101* knock-out cells in a *Cdkn2a* heterozygous background (*Tsg101*^{fl/fl} *Cdkn2a*^{+/-}) or *Tsg101* heterozygous mutants carrying two *Cdkn2a* null alleles (*Tsg101*^{fl/+} *Cdkn2a*^{-/-}) were used as controls. The infection of these control cell lines with the pBabe retroviral control vector and puromycin selection had also no influence on cell survival (data not shown). As expected, the expression of Cre also had no effect on cell proliferation and cell survival of MEFs with a *Tsg101*-floxed heterozygous mutation whether cells express p19^{Arf} or not. However, the ablation of *Tsg101* in p19^{Arf}-deficient cells using the pBabe-Cre construct (*Tsg101*^{fl/fl} *Cdkn2a*^{-/-} pBabe-Cre) resulted in cell death shortly after excision of *Tsg101*, and the phenotype was comparable to those observed in *Tsg101/p53* and *Tsg101/p21* double mutants (data not shown). This observation suggested that none of the genes of the p19^{Arf}-Mdm2-p53-p21^{Cip1} circuit alone or in combination with p16^{Ink4a} seem to be required for the initiation of cell death of *Tsg101*-deficient MEFs.

Our previous studies showed that the conditional deletion of the *Tsg101* gene did not negatively affect the steady-state level of Mdm2 in primary fibroblasts (12). Similarly, the steady-state level of Mdm2 was not reduced but was slightly elevated in *Cdkn2a*^{+/-} or *Cdkn2a*^{-/-} MEFs lacking *Tsg101* expression (Fig. 5B). We performed a flow cytometric analysis to study whether the loss of *Tsg101* function was able to restore the G₁ arrest in p19^{Arf}-deficient cells before they died (Fig. 5C). In addition, we used p19^{Arf}-deficient cells (*Tsg101*^{fl/fl} *Cdkn2a*^{-/-}) deprived of essential growth factors (0.1% fetal bovine serum) as a positive control to monitor the accumulation of cells at G₁ and the relative decrease in the number of cells at the S phase of the cell cycle. Cells deficient in *Tsg101* and p19^{Arf} did not arrest at G₁ and had approximately the same relative number

² C. Eischen (University of Nebraska Medical Center), J. Weber (Washington University), and G. Zambetti (St. Jude's Children's Research Hospital), personal communications.

of cells in S phase as MEFs lacking Tsg101 in addition to p53 and/or p21^{Cip1} (see Fig. 2A and 3B; $p > 0.05$).

To verify these findings in a different experimental setting, we immortalized *Tsg101*^{f/f} MEFs and their wild type controls by inhibiting the expression of p19^{Arf} through overexpression of the T-box protein 2 (data not shown). T-box protein 2 is a known transcriptional repressor for the mouse and human *CDKN2A(ARF)* promoters (34). MEFs expressing T-box protein 2 exhibited reduced levels of p19^{Arf} and p21^{Cip1} at passages seven and eight. After Cre-mediated recombination of the floxed *Tsg101* locus, these immortalized cells died within 7 days post-infection. The steady-state levels of Mdm2 did not decrease, and p21^{Cip1} protein expression levels remained low and unchanged in the Tsg101 knock-out cells lacking functional p19^{Arf} (data not shown). In summary, these results are consistent with our findings in Tsg101 knock-out cells with a targeted deletion of the entire *Cdkn2a* locus. Therefore, Tsg101 deficiency seemed to not alter Mdm2 function in a biologically relevant manner that would influence the Mdm2/p53 negative feedback loop.

Transforming Mutations during Neoplastic Transformation Do Not Affect the Important Role of Tsg101 for Cell Survival—Li and Cohen (16) report that a functional knock-out of *Tsg101* using a conventional antisense approach and inducible overexpression of *Tsg101* resulted in reversible neoplastic transformation of mouse 3T3 fibroblasts. Our published observations using a site-directed, targeted knock-out in animal models and derived cell lines suggested that a loss-of-function of *Tsg101* is insufficient to trigger neoplastic transformation *in vitro* and *in vivo* (12, 13). Because *Tsg101* seems not to be a primary tumor suppressor, we hypothesized that this gene might function as a modifier for neoplastic transformation (12). Null or inactivating mutations in p19^{Arf}, p53, or p21^{Cip1} are able to partially rescue the complex phenotype caused by Tsg101 deficiency (*i.e.* the G₁/S progression), and it is, therefore, logical to test whether additional mutations during neoplastic transformation have an impact on the crucial role of Tsg101 as a survival factor in tumorigenic cells. Fig. 6A illustrates our experimental design to address this issue. We previously described the generation of immortalized *Tsg101*^{f/f} MEFs using a standard 3T3 protocol (12). These cells carry an E255D mutation in the DNA binding domain of p53, and consequently, the p21^{Cip1} protein is not expressed. The deletion of *Tsg101* in this immortalized cell line resulted in instant cell death (12). The G₁/S transition is, however, restored due to the lack of p21^{Cip1}. Hence, the phenotype of these cells relating to cell cycle regulation and cell death is equivalent to double-mutant MEFs with targeted deletions of *p53* and *Tsg101*.³ Immortal *Tsg101*^{f/f} 3T3 cells were passaged numerous times, infected with a pBabe-hygro retrovirus, and selected for hygromycin resistance. Next, 8×10^5 cells were injected subcutaneously into athymic nude mice (NCr strain) to select mutants that were able to grow *in vivo* and to form solid tumors. Tumorigenesis was observed in all animals after a medium latency of 3–5 weeks (Fig. 6B). Histopathologically, these lesions were well vascularized, and neoplastic cells invaded into adjacent normal tissues such as the fat pad of the thoracic mammary gland (Fig. 6C). Tumor cells were explanted and grown in culture in hygromycin-containing media to remove all nontumorigenic cell types such as endothelial cells, epithelial cells, and tumor-associated fibroblasts of the host. Next, neoplastic cells were infected with pBabe-Cre to excise both floxed copies of *Tsg101*. Infection of these cells with the control vector pBabe and selection with puromycin had little effect on their viability, whereas the deletion of *Tsg101*

resulted in growth arrest and cell death (Fig. 6, D and E). These data suggested that additional transforming mutations during neoplastic transformation did not rescue the lethal phenotype caused by Tsg101 deficiency, and therefore, Tsg101 might be important for the survival of both normal and neoplastic cells.

DISCUSSION

Tsg101 Deficiency and Cellular Stress Response—We previously reported that Tsg101 is indispensable for the survival of cells *in vitro* and *in vivo* (12, 13). The collective data presented in this new report demonstrated that targeted mutations to abolish the function of p19^{Arf}, p16^{Ink4a}, p53, and p21^{Cip1} were insufficient to maintain the survival of Tsg101 knock-out cells. These observations suggest that the p19^{Arf}-Mdm2-p53 tumor surveillance or stress response pathway does not control cellular mechanisms that lead to the death of Tsg101-deficient cells. These results are in full agreement with our previous studies, in which we have demonstrated that the functional inhibition of p53, through expression of E6 or a spontaneous mutation within the DNA binding domain of p53, had no effect on the viability of Tsg101 knock-out cells (12). By analyzing conventional knock-out mice that die very early *in utero*, Ruland *et al.* (11) established that Tsg101 is required for cell proliferation and that p53 is a crucial player for the replication defect of Tsg101-deficient cells. Using our conditional knock-out model, we established that the deletion of *Tsg101* resulted in a sustained cell cycle arrest at the G₁ checkpoint before these cells died (12). Based on the findings by Ruland *et al.* (11) and due to the fact that Tsg101 conditional knock-out cells lacked active Cdk2, we hypothesized that the p53 and its downstream mediator p21^{Cip1} may at least in part contribute to the complex phenotype. We proposed that cell cycle arrest and cell death were controlled by separate, p53-dependent and p53-independent, pathways (13). In this report we show that, in contrast to mechanisms leading to cell death, the G₁ arrest in Tsg101-deficient cells is dependent upon expression of functional p53. We also determined that p21^{Cip1} is an integral part of this phenotype. Recent reports suggested a potential role for Tsg101 as a dominant-negative regulator for Mdm2 and p21^{Cip1} protein turnover (17, 18). Evidently, Tsg101 cannot serve simultaneously as a stabilizer for both regulators of the cell cycle. According to the model by Li *et al.* (17) and Ruland *et al.* (11), functional levels of p21^{Cip1} were required to mediate the cell cycle arrest. Tsg101 deficiency leads to destabilization of Mdm2, accumulation of active p53, and subsequently, the transcriptional activation of p21^{Cip1}. If Tsg101 is required for p21 stability or the recruitment of p21 to the cyclin-Cdk2 complex as proposed by Oh *et al.* (18), then the combined effects of a Tsg101 knock-out on Mdm2 and p21^{Cip1} should nullify each other.

The overexpression of Tsg101 or its conditional deletion results in a G₁ arrest and cell death, suggesting that the level of Tsg101 protein in a given cell is tightly regulated and that a high or low amount of Tsg101 outside a narrow range cannot be tolerated (9, 10, 12, 13, 35, 36). Interestingly, p21^{Cip1} seems to be a key player for the G₁ arrests not only in the overexpression model (18) but also the conditional knock-out. As demonstrated in this report, a knock-out of the *Cdkn1a* gene (p21^{Cip1}) eliminated the G₁ arrest and restored the G₁/S progression of Tsg101-deficient cells. Based on the fact that p21^{Cip1} levels did not change in the Tsg101 conditional knock-out (Ref. 12 and Figs. 2B, second and third lanes, and 3A, fifth and sixth lanes), we presume that Tsg101 deficiency leads directly to cellular conditions where p21 ability to bind to the Cdk2 is elevated. The pathways that link the function of Tsg101 and p21^{Cip1} need to be identified. There is increasing evidence in the recent literature that Tsg101 is essential for endosomal trafficking

³ A. Krempler and K.-U. Wagner, unpublished information.

and other cellular processes in addition to the regulation of ubiquitination (5–8). It is, therefore, very likely that Tsg101 deficiency or its overexpression will cause a severe negative impact on a number of biological processes. Those might subsequently trigger the activation of cellular stress responses such as the recruitment of p21^{Cip1} to the Cdk2 complex to arrest cells in the G₁ phase. Ruland *et al.* (11) suggest that p53 is the upstream regulator of p21^{Cip1}, which mediates the cell cycle arrest in Tsg101-deficient cells. Based on data by Li *et al.* (17), the authors proposed a direct interaction of Tsg101 with members of the p53 stress response pathway, in particular Mdm2. In the conditional knock-out model we were, however, unable to detect higher levels of p53 or changes in the p53 transcriptional activation of p21^{Cip1} (12). Moreover, we questioned the suggested role of Tsg101 as a positive regulator for Mdm2 function based on the evidence that *Tsg101/p53* double mutant mice still die early *in utero* (11), whereas p53 deficiency results in a complete rescue of embryonic lethality in *Mdm2* knock-out mice (31, 32). To rigorously examine the proposed function of Tsg101 as a stabilizer for Mdm2, we generated cells lacking Tsg101 and p19^{Arf}. The functional inhibition of p19^{Arf} leads to increased Mdm2-mediated sequestration of p53 and immortalization (20). We utilized the complete deletion of the *Cdkn2a* locus and the T-box protein 2-mediated transcriptional repression of p19^{Arf} as two alternative approaches to immortalize MEFs carrying two conditional *Tsg101* knock-out alleles. In both model systems the deletion of *Tsg101* had no effect on Mdm2 steady-state levels and function. In particular, the p53-mediated transcriptional activation of p21^{Cip1} and induction of a G₁ arrest was not restored. In addition, we were unable to co-precipitate Mdm2 with endogenous Tsg101 or transgenic hemagglutinin-tagged Tsg101 expressed at near physiological levels,⁴ suggesting that under normal conditions these proteins show weak or no interaction. In summary, results obtained from three double knock-out models (*Tsg101/p53*, *Tsg101/p21*, and *Tsg101/p19^{Arf}*) suggest that a direct interaction of Tsg101 with members of the Mdm2-p53 circuit is unlikely. We propose that the activation of a p21^{Cip1}-mediated G₁ arrest might be an indirect effect of Tsg101 deficiency on other cellular processes that subsequently trigger stress response pathways.

Normal Tsg101 Function Is Crucial for the Survival of Primary, Immortalized, and Neoplastic Cells—The role of the *TSG101* gene as a tumor suppressor or oncogene in human malignancies is still controversial. After cloning the mouse *Tsg101* locus and revising the human *TSG101* gene structure, we found that many of the previously described aberrant splice variants in human malignancies represent alternative splice forms that originate solely from exon skipping (35, 37). Although the transforming capability of *Tsg101* as an oncogene has not been examined in an animal model to date, we and others could show that neither haplo-insufficiency of *Tsg101* (11, 13) nor the deletion of both *Tsg101* alleles in selected cell types of conditional knock-out mice resulted in tumorigenesis (12, 13). In contrast to a previous report (16), we demonstrated that a null mutation of *Tsg101* is not an initiating event for neoplastic transformation of primary cells (12, 13). Therefore, we hypothesized that this gene might function as a modifier for tumorigenesis (12). This earlier assumption was based on the fact that the role of Tsg101 in neoplastic transformation was initially established in immortalized 3T3 cells (16). In preliminary studies we could show that these cells lacked expression of other important tumor suppressor proteins, in particular p19^{Arf} and

p16^{Ink4a} (see Fig. 5). As discussed above, Tsg101 was also suggested to be a crucial regulator of the Mdm2-p53 circuit and p21^{Cip1}. In contrast to our earlier working hypothesis, we can demonstrate in this study that none of the immortalizing mutations (*i.e.* knock-out of p19^{Arf}, p16^{Ink4a}, p53, and/or p21^{Cip1}) or additional sporadic mutations causing neoplastic transformation and tumorigenesis had an impact on the important role of Tsg101 for cell survival. In agreement with these findings in primary cell cultures, the deletion of the *p53* gene in *Tsg101* mammary-specific knock-out mice (WAP-Cre *Tsg101^{fl/fl} p53^{-/-}*) did not rescue the survival of Tsg101-deficient epithelial cells. More importantly, haplo-insufficiency of *p53* in the Tsg101 somatic knock-out model did not cause mammary tumorigenesis in 12-month-old animals. Unlike in our *Brcal* mammary-specific knock-out mice (38), the deletion of *Tsg101* in *p53* heterozygous mutants (WAP-Cre *Tsg101^{fl/fl} p53^{+/-}*) seemed not to cause genomic alterations that lead to the loss of heterozygosity of the wild type *p53* allele and neoplastic transformation.⁴ Mammary tumors were also not observed in females carrying two mutant *p53* alleles (WAP-Cre *Tsg101^{fl/fl} p53^{-/-}*). Like *p53* single knock-out mice, they succumbed to lymphoma after a latency period of approximately six months. Finally, we have confirmed the importance of Tsg101 as a survival factor for tumorigenic cells in a murine cancer model that carries a mouse mammary tumor virus (MMTV)-driven *Her2/neu* oncogene (39) in addition to the somatic knock-out of *Tsg101* (MMTV-neu WAP-Cre *Tsg101^{fl/fl}*). In this animal model the deletion of *Tsg101* did not accelerate but delayed neoplastic transformation (40). In conclusion to the data presented in this study, we propose that Tsg101 is an important factor for cell survival in normal, immortalized, and neoplastic cells. Because Tsg101 is essential for cell survival, the ability to target tumor cells and inhibit Tsg101 function should result in the death of neoplastic cells regardless of the functional inactivation of the Mdm2-p53 circuit or sporadic mutations in the p53, p19^{Arf}, or p16^{Ink4a} tumor susceptibility loci.

Acknowledgments—We thank Malinda D. Henry for technical assistance and animal husbandry. We also thank Charles Kuszynski and Linda Wilkie of the University of Nebraska Medical Center Flow Cytometry Facility for the cell cycle analysis and Joerg Rahnenfuhrer (University of California, Berkeley) for help in the evaluation of the MTT assay. We are grateful to Dr. Alan Diehl (University of Pennsylvania) for reading this manuscript and valuable suggestions.

REFERENCES

- Koonin, E. V., and Abagyan, R. A. (1997) *Nat. Genet.* **16**, 330–331
- Ponting, C. P., Cai, Y. D., and Bork, P. (1997) *J. Mol. Med.* **75**, 467–469
- Watanabe, M., Yanagi, Y., Masuhiro, Y., Yano, T., Yoshikawa, H., Yanagisawa, J., and Kato, S. (1998) *Biochem. Biophys. Res. Commun.* **245**, 900–905
- Hittelman, A. B., Burakov, D., Iniguez-Lluhi, J. A., Freedman, L. P., and Garabedian, M. J. (1999) *EMBO J.* **18**, 5380–5388
- Babst, M., Odorizzi, G., Estepa, E. J., and Emr, S. D. (2000) *Traffic* **1**, 248–258
- Garrus, J. E., von Schwedler, U. K., Pornillos, O. W., Morham, S. G., Zavitz, K. H., Wang, H. E., Wettstein, D. A., Stray, K. M., Cote, M., Rich, R. L., Myszkka, D. G., and Sundquist, W. I. (2001) *Cell* **107**, 55–65
- Lu, Q., Hope, L. W., Brasch, M., Reinhard, C., and Cohen, S. N. (2003) *Proc. Natl. Acad. Sci. U. S. A.* **100**, 7626–7631
- Bache, K. G., Brech, A., Mehlum, A., and Stenmark, H. (2003) *J. Cell Biol.* **162**, 435–442
- Zhong, Q., Chen, Y., Jones, D., and Lee, W. H. (1998) *Cancer Res.* **58**, 2699–2702
- Xie, W., Li, L., and Cohen, S. N. (1998) *Proc. Natl. Acad. Sci. U. S. A.* **95**, 1595–1600
- Ruland, J., Sirard, C., Elia, A., MacPherson, D., Wakeham, A., Li, L., Luis, D. L. P., Cohen, S. N., and Mak, T. W. (2001) *Proc. Natl. Acad. Sci. U. S. A.* **98**, 1859–1864
- Krempler, A., Henry, M. D., Triplett, A. A., and Wagner, K. U. (2002) *J. Biol. Chem.* **277**, 43216–43223
- Wagner, K. U., Krempler, A., Qi, Y., Park, K., Henry, M. D., Triplett, A. A., Riedlinger, G., Rucker, E. B., III, and Hennighausen, L. (2003) *Mol. Cell. Biol.* **23**, 150–162
- Maucuer, A., Camonis, J. H., and Sobel, A. (1995) *Proc. Natl. Acad. Sci. U. S. A.* **92**, 3100–3104
- Sun, Z., Pan, J., Hope, W. X., Cohen, S. N., and Balk, S. P. (1999) *Cancer* **86**,

⁴ A. Krempler, A. A. Triplett, and K.-U. Wagner, unpublished information.

- 689–696
16. Li, L., and Cohen, S. N. (1996) *Cell* **85**, 319–329
 17. Li, L., Liao, J., Ruland, J., Mak, T. W., and Cohen, S. N. (2001) *Proc. Natl. Acad. Sci. U. S. A.* **98**, 1619–1624
 18. Oh, H., Mammucari, C., Nenci, A., Cabodi, S., Cohen, S. N., and Dotto, G. P. (2002) *Proc. Natl. Acad. Sci. U. S. A.* **99**, 5430–5435
 19. Li, L., Li, X., Francke, U., and Cohen, S. N. (1997) *Cell* **88**, 143–154
 20. Sherr, C. J. (1998) *Genes Dev.* **12**, 2984–2991
 21. Wagner, K. U., Wall, R. J., St-Onge, L., Gruss, P., Wynshaw-Boris, A., Garrett, L., Li, M., Furth, P. A., and Hennighausen, L. (1997) *Nucleic Acids Res.* **25**, 4323–4330
 22. Donehower, L. A., Harvey, M., Slagle, B. L., McArthur, M. J., Montgomery, C. A., Jr., Butel, J. S., and Bradley, A. (1992) *Nature* **356**, 215–221
 23. Brugarolas, J., Chandrasekaran, C., Gordon, J. I., Beach, D., Jacks, T., and Hannon, G. J. (1995) *Nature* **377**, 552–557
 24. Serrano, M., Lee, H., Chin, L., Cordon-Cardo, C., Beach, D., and DePinho, R. A. (1996) *Cell* **85**, 27–37
 25. Wagner, K. U., Boulanger, C. A., Henry, M. D., Sgagias, M., Hennighausen, L., and Smith, G. H. (2002) *Development* **129**, 1377–1386
 26. Telford, W. G., King, L. E., and Fraker, P. J. (1991) *Cell Prolif.* **24**, 447–459
 27. van de Loosdrecht, A. A., Beelen, R. H., Ossenkuppele, G. J., Broekhoven, M. G., and Langenhuijsen, M. M. (1994) *J. Immunol. Methods* **174**, 311–320
 28. el Deiry, W. S., Tokino, T., Velculescu, V. E., Levy, D. B., Parsons, R., Trent, J. M., Lin, D., Mercer, W. E., Kinzler, K. W., and Vogelstein, B. (1993) *Cell* **75**, 817–825
 29. Xiong, Y., Hannon, G. J., Zhang, H., Casso, D., Kobayashi, R., and Beach, D. (1993) *Nature* **366**, 701–704
 30. Sherr, C. J., and Roberts, J. M. (1999) *Genes Dev.* **13**, 1501–1512
 31. Jones, S. N., Roe, A. E., Donehower, L. A., and Bradley, A. (1995) *Nature* **378**, 206–208
 32. Montes de Oca, L. R., Wagner, D. S., and Lozano, G. (1995) *Nature* **378**, 203–206
 33. Weber, J. D., Jeffers, J. R., Reh, J. E., Randle, D. H., Lozano, G., Roussel, M. F., Sherr, C. J., and Zambetti, G. P. (2000) *Genes Dev.* **14**, 2358–2365
 34. Jacobs, J. J., Keblusek, P., Robanus-Maandag, E., Kristel, P., Lingbeek, M., Nederlof, P. M., van Welsem, T., van de Vijver, M. J., Koh, E. Y., Daley, G. Q., and van Lohuizen, M. (2000) *Nat. Genet.* **26**, 291–299
 35. Wagner, K. U., Dierisseau, P., Rucker, E. B., Robinson, G. W., and Hennighausen, L. (1998) *Oncogene* **17**, 2761–2770
 36. Feng, G. H., Lih, C. J., and Cohen, S. N. (2000) *Cancer Res.* **60**, 1736–1741
 37. Wagner, K. U., Dierisseau, P., and Hennighausen, L. (1999) *Cytogenet. Cell Genet.* **84**, 87–88
 38. Xu, X., Wagner, K. U., Larson, D., Weaver, Z., Li, C., Ried, T., Hennighausen, L., Wynshaw-Boris, A., and Deng, C. X. (1999) *Nat. Genet.* **22**, 37–43
 39. Guy, C. T., Webster, M. A., Schaller, M., Parsons, T. J., Cardiff, R. D., and Muller, W. J. (1992) *Proc. Natl. Acad. Sci. U. S. A.* **89**, 10578–10582
 40. M. D. Henry Henry, M. D., Triplett, A. A., Oh, K.B., Smith, G. H., and Wagner, K.-U. (2004) *Oncogene*, in press